



Claritin-D 24 Hour Tablets

Drug Facts

Active ingredients (in each tablet)	Purpose
Loratadine 10 mg.....	Antihistamine
Pseudoephedrine sulfate 240 mg.....	Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing ■ itchy, watery eyes
 - runny nose ■ itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease
- high blood pressure ■ diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not divide, crush, chew or dissolve the tablet	
adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **each tablet contains:** calcium 25 mg
- safety sealed: do not use if the individual blister unit imprinted with Claritin-D® 24 hour is open or torn
- store between 20°C to 25°C (68°F to 77°F)
- protect from light and store in a dry place

Inactive ingredients carnauba wax, dibasic calcium phosphate dihydrate, ethylcellulose, hydroxypropyl cellulose, hypromellose, magnesium stearate, pharmaceutical ink, polyethylene glycol, povidone, silicon dioxide, sucrose, titanium dioxide

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com



Claritin-D 24 Hour Tablets