



# Claritin-D 12 Hour Tablets

## Drug Facts

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Loratadine 5 mg.....	Antihistamine
Pseudoephedrine sulfate 120 mg.....	Nasal Decongestant

### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - itchy, watery eyes
  - runny nose
  - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

### Warnings

#### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

#### When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

<b>Directions</b>	
■ do not divide, crush, chew or dissolve the tablet	
adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

### Other information

- each tablet contains: calcium 30 mg
- safety sealed: do not use if the individual blister unit imprinted with Claritin-D® 12 hour is open or torn
- store between 20°C to 25°C (68°F to 77°F)
- keep in a dry place

**Inactive ingredients** croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

### Questions or comments?

1-800-CLARITIN (1-800-252-7484) or [www.claritin.com](http://www.claritin.com)